

AD-A089 011

ROYAL AUSTRALIAN NAVY RESEARCH LAB EDGECLIFF
A TIME SERIES OF OCEANOGRAPHIC SECTIONS ACROSS THE EQUATOR AT 9--ETC(U)
NOV 79 P J MULHEARN, R J EDWARDS

F/G 8/10

UNCLASSIFIED

RANRL-TN-2/79

NL

1-1

2-1

3-1

4-1

5-1

6-1

7-1

8-1

9-1

10-1

11-1

12-1

13-1

14-1

15-1

16-1

17-1

18-1

19-1

20-1

21-1

22-1

23-1

24-1

25-1

26-1

27-1

28-1

29-1

30-1

31-1

32-1

33-1

34-1

35-1

36-1

37-1

38-1

39-1

40-1

41-1

42-1

43-1

44-1

45-1

46-1

47-1

48-1

49-1

50-1

51-1

52-1

53-1

54-1

55-1

56-1

57-1

58-1

59-1

60-1

61-1

62-1

63-1

64-1

65-1

66-1

67-1

68-1

69-1

70-1

71-1

72-1

73-1

74-1

75-1

76-1

77-1

78-1

79-1

80-1

81-1

82-1

83-1

84-1

85-1

86-1

87-1

88-1

89-1

90-1

91-1

92-1

93-1

94-1

95-1

96-1

97-1

98-1

99-1

100-1

101-1

102-1

103-1

104-1

105-1

106-1

107-1

108-1

109-1

110-1

111-1

112-1

113-1

114-1

115-1

116-1

117-1

118-1

119-1

120-1

121-1

122-1

123-1

124-1

125-1

126-1

127-1

128-1

129-1

130-1

131-1

132-1

133-1

134-1

135-1

136-1

137-1

138-1

139-1

140-1

141-1

142-1

143-1

144-1

145-1

146-1

147-1

148-1

149-1

150-1

151-1

152-1

153-1

154-1

155-1

156-1

157-1

158-1

159-1

160-1

161-1

162-1

163-1

164-1

165-1

166-1

167-1

168-1

169-1

170-1

171-1

172-1

173-1

174-1

175-1

176-1

177-1

178-1

179-1

180-1

181-1

182-1

183-1

184-1

185-1

186-1

187-1

188-1

189-1

190-1

191-1

192-1

193-1

194-1

195-1

196-1

197-1

198-1

199-1

200-1

201-1

202-1

203-1

204-1

205-1

206-1

207-1

208-1

209-1

210-1

211-1

212-1

213-1

214-1

215-1

216-1

217-1

218-1

219-1

220-1

221-1

222-1

223-1

224-1

225-1

226-1

227-1

228-1

229-1

230-1

231-1

232-1

233-1

234-1

235-1

236-1

237-1

238-1

239-1

240-1

241-1

242-1

243-1

244-1

245-1

246-1

247-1

248-1

249-1

250-1

251-1

252-1

253-1

254-1

255-1

256-1

257-1

258-1

259-1

260-1

261-1

262-1

263-1

264-1

265-1

266-1

267-1

268-1

269-1

270-1

271-1

272-1

273-1

274-1

275-1

276-1

277-1

278-1

279-1

280-1

281-1

282-1

283-1

284-1

285-1

286-1

287-1

288-1

289-1

290-1

291-1

292-1

293-1

294-1

295-1

296-1

297-1

298-1

299-1

300-1

301-1

302-1

303-1


304-1

305-1

306-1

307-1

308-1</



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS 1963 A

LEVEL II
UNCLASSIFIED

AR Number: AR-001-499



DEPARTMENT OF DEFENCE
DEFENCE SCIENCE AND TECHNOLOGY ORGANISATION
R.A.N. RESEARCH LABORATORY
EDGECLIFF, N.S.W.

RANRL TECHNICAL NOTE No. 2/79

© COMMONWEALTH OF AUSTRALIA 1979

**A TIME SERIES OF OCEANOGRAPHIC SECTIONS
ACROSS THE EQUATOR AT 92°E IN THE
EASTERN INDIAN OCEAN**

BY
**P. J. MULHEARN, RANRL and R. J. EDWARDS, CSIRO Div of
FISHERIES OCEANOGRAPHY**



APPROVED FOR PUBLIC RELEASE

THE UNITED STATES NATIONAL
TECHNICAL INFORMATION SERVICE
IS AUTHORISED TO
REPRODUCE AND SELL THIS REPORT

COPY No. 14

NOVEMBER 1979

UNCLASSIFIED

80 9 11 046

AD A089011

FILE COPY

DTIC
ELECTED
SEP 12 1980
C

406 607

DOCUMENT CONTROL DATA SHEET

Security classification of this page:

1. <u>DOCUMENT NUMBERS:</u>		2. <u>SECURITY CLASSIFICATION:</u>	
a. AR Number AR-001-499		a. Complete document: UNCLASSIFIED	
b. Document Series and Number: RANRL Technical Note		b. Title in isolation: "	
c. Report Number: 2/79		c. Summary in Isolation: "	
3. <u>TITLE</u> A time series of oceanographic sections across the equator at 92°E in the eastern Indian Ocean.			
4. <u>PERSONAL AUTHOR(S):</u> P.J. Mulhearn, R.J. Edwards		5. <u>DOCUMENT DATE:</u> Nov 1979	
7. <u>CORPORATE AUTHOR(S):</u> RAN Research Laboratory PO Box 706 Darlinghurst NSW 2010		6. <u>TYPE OF REPORT AND PERIOD COVERED:</u> Progress May and June 1979	
		8. <u>REFERENCE NUMBERS:</u> a. Task: NAV Task 76/059 b. Sponsoring Agency: NSA Approved: <i>W.F. Hunter</i> SSRL	
10. <u>IMPRINT</u> (Publishing establishment) RAN Research Laboratory		9. <u>COST CODE</u>	
		11. <u>COMPUTER PROGRAM(S):</u> (Title(s) and Language(s)) NIL	
12. <u>RELEASE LIMITATIONS</u> (of the document): (in addition to the Distribution List) AUSTRALIA: approved for public release			
12-0. <u>OVERSEAS:</u> N.O. <input type="checkbox"/> P.R. <input checked="" type="checkbox"/> 1 A <input type="checkbox"/> B <input type="checkbox"/> C <input type="checkbox"/> D <input type="checkbox"/> E <input type="checkbox"/>			
13. <u>ANNOUNCEMENT LIMITATIONS</u> (of the information on this page): No limitation			
14. <u>DESCRIPTORS:</u> Physical Oceanography Oceanographic Surveys		15. <u>COSATI CODES:</u> 0810	
16. <u>SUMMARY/ABSTRACT:</u>			

> A series of oceanographic sections across the equator in May and June 1979 were carried out by RANRL and CSIRO at 92°E in the eastern Indian Ocean. Marked reversals in the surface currents were found which suggest the presence of either equatorially trapped waves or waves caused by the Ninety East Ridge. The XBT sections and dynamic height contours support the observed current reversals.

DISTRIBUTION LIST

	<u>Copy No</u>
Chief Defence Scientist	1
Deputy Chief Defence Scientist	2
Superintendent Scientific and Technical Programme	3
Controller, Service Laboratories and Trails Division	4
Joint Intelligence Organisation (DDSTI)	5
Defence Library, Campbell Park	6
Defence Information Services Branch - Microfiche	7
- release to US - DDC	8 - 19
- release to UK - DRIC	20
- release to Canada - DSIS	21
- release to NZ - DoD	22
Director, Naval Meteorological and Oceanographic Services	23
Deputy Director, Naval Meteorological and Oceanographic Services	24
Mr D. Rochford, CSIRO Division of Fisheries and Oceanography	25
Mr R. Edwards, CSIRO Division of Fisheries and Oceanography	26
Library, RAN Research Laboratory	27 - 38
Library, CSIRO Division of Fisheries and Oceanography	39 - 49
Dr L. Mysak, University of British Columbia, Vancouver, Canada	50
Dr J. Swallow, Present SLOR Indian Ocean	51
Dr M. Fleux, Laboratoire d'Océanographie Physique, Paris, France	52
Dr S.G.H. Philander, NOAA/GFDL, Princeton University, Princeton, New Jersey 08540, USA	53

CONTENTS

	<u>Page No</u>
1. Introduction	1
2. Methods	2
3. Results	2
3.1 XBT Sections	2
3.2 Surface Currents	2
3.3 Dynamic Heights	3
3.4 Satellite-tracked buoys	3
4. Conclusion	3
5. Acknowledgements	3
6. References	3




Fig. 1. Cruise tracks (Dashed line shows deviation from 92°E on RANRL's return leg).

1. Introduction

In May and June 1979 the R.A.N. Research Laboratory and the CSIRO Division of Fisheries and Oceanography carried out a series of North-South oceanographic sections across the equator at 92°E, in the eastern Indian Ocean. The experiments were part of the oceanographic programme of the First GARP Global Experiment (FGGE) and occurred in its Second Special Observing Period. (See GARP Newsletters). A primary aim of FGGE's oceanographic programme was to obtain a better understanding of equatorial current systems. In the Indian Ocean most interest was centred on the western half and the Somali Current, so that the results discussed here represent one of the few data sets for the eastern half of the Indian Ocean at that time. Other data was obtained from U.S.S. Wilkes and some may have been obtained by a polygon of five Russian ships which traversed along the equator before and after Special Observing Period II.

The coverage and dates of the RANRL and CSIRO cruises, all using HMAS Diamantina are set out below.

<u>Dates</u>	<u>Latitude and Longitude</u>	<u>Institution</u>
12-15 May	5°N, 92°E to 3°S, 92°E	RANRL
15-17 May	3°S, 92°E to ½°N, 92°E	RANRL
17-18 May	½°N, 92°E to 6°N, 95°E	RANRL
28 May - 1 June	5°N, 92°E to 5°S, 92°E	CSIRO
1-5 June	5°S, 92°E to 5°N, 92°E	CSIRO

RANRL only went as far south as 3°S because it was taking longer time-series measurements en-route. It also had to leave the 92°E line on 17 May and head for Singapore because of a fire on board. Cruise tracks are shown on Figure 1.

Some data were taken by RANRL which were not taken by CSIRO and vice-versa, because of the slightly different aims of the two cruises. The data presented here are those which were measured by both institutions and hence allow a time series development of certain oceanographic parameters to be presented. Data to be presented are

- (1) Temperature cross-sections from XBT's
- (2) Surface currents
- (3) Dynamic heights
- (4) Satellite-tracked buoy paths

Some previous work on the eastern equatorial Indian Ocean is contained in works by Wyrski (1973), Taft and Knauss (1967), Eriksen (1979), O'Brien and Hurlburt (1974) and Philander (1973). From climatological data Wyrski showed the existence of an eastward equatorial jet in the




Fig. 2(a). XBT section on south-going RANRL leg 12/5/79-15/5/79.

NOV. 79




Fig. 2(b). XBT section on north-going RANRL leg 15/5/79-18/5/79.

NOV. 79

monsoon changeover months of May and October. O'Brien and Hurlburt obtained such a jet in a simple numerical model which imposed a uniform eastward wind over the ocean basin - such eastward winds being predominant in May and October. Taft and Knauss at 53°E, found a surface eastward equatorial current in May in the course of one of their cruises. Data from other times usually show a westward flow at the surface with a subsurface eastwards jet at the equator. Philander (1973) provides a broad review of the equatorial current systems.

2. Methods

The Temperature sections to be presented were all obtained with standard XBT equipment. The surface currents from RANRL's cruise were obtained when the ship was drifting on station and obtained four or more sat-nav fixes. Surface currents from CSIRO's cruise were obtained by the differences between dead-reckoning and sat-nav fixes.

For RANRL's cruise dynamic heights were obtained using temperatures from XBT's and salinities from shallow Niskin casts and XSTD probe data. (XSTD = expendable salinity - temperature depth). No reversing thermometers were attached to the Niskin bottles but the true depths of the salinity samples were obtained from the angle of the hydrographic wire and a comparison of the shapes of salinity profiles from the Niskin bottles and the XSTD probes. CSIRO obtained their dynamic heights from Nansen and Niskin cast data by standard means. All salinities were measured on board using an inductive salinometer (Brown and Hamon, 1961).

Satellite-tracked buoys were deployed by CSIRO on 1 June at 0° and 3 June at 1°N, both on 92°E. These were of the torpedo shape described in Cresswell and others (1978) and were drogued at 20 m.

3. Results

3.1 XBT Sections

The four XBT sections in Figures 2(a) to (d) are rather featureless, the isotherms in most cases being almost horizontal, except that for the RANRL sections there is a tendency for isotherms near the base of the mixed layer to converge near the equator. This convergence suggests that the surface flow is to the east relative to the water below the thermocline. The section of Fig 2(b) shows more features especially along the track from ½°N, 92°E to 5°N, 94°20'E. The time series at 0° shown in Fig 2(e) shows very little change over a ten hour period.

3.2 Surface Currents

Ship's sets are available at the positions shown on Figures 3(a) and (b). They are available for the southward leg of RANRL's cruise from 12-15 May and on 16 May at 0°, during the northwards leg. On CSIRO's cruise they are available only for the southward leg from 28 May to 1 June, because the sat-nav failed on the way back north.




Fig. 2(c). XBT section on south-going CSIRO leg 28/5/79-1/6/79.




Fig. 2(d) XBT section on north-going CSIRO leg 1/6/79-5/6/79.

The reversal of the currents (some of up to 2 knots magnitude) in the two weeks between the cruises is a striking feature of these results. From 12-15 May currents were to the east while they went to the west in the 28 May - 1 June period. Winds were light and variable throughout this period and a calm prevailed for most of the RANRL legs.

3.3 Dynamic Heights

Profiles of dynamic height of the sea-surface relative to 300 m, taken for the 3 periods 12-15 May, 28 May - 1 June and 1-5 June are presented in Figs 4(a) to (c). Dynamic heights are larger and show more variation for the period 12-15 May than for the other two legs. Geostrophic currents calculated from these dynamic topographies appear to be of the correct order north of $1\frac{1}{2}^{\circ}\text{N}$ and south of $1\frac{1}{2}^{\circ}\text{S}$ but the amount of reliable data on currents available by this method is very limited. The change in slope of the dynamic height between 2°N and 4°N from RANRL's leg of 12-15 May and CSIRO's leg of 28 May - 1 June supports the surface current measurements. Overall there is a tendency for the range of dynamic height to decrease with time.

The dynamic height relative to 1000 dbars is shown in Fig 4(d) for the period 28 May - 1 June.

3.4 Satellite-tracked Buoys

Satellite-tracked buoys were deployed on 1 June at 0°N , 92°E and on 3 June at 1°N , 92°E . The tracks of these buoys follow each other closely for the next 80 days at least. Initially they drifted westwards for 12 to 14 days, supporting the ships set data, then turned back towards the east and continued towards the east, tending southwards for another 78 days before turning towards the west once more. The final journey to the west may be due to capture by the South Equatorial Current, but the earlier reversing motion revealed by both the buoy tracks and the ship's sets may be caused by equatorially trapped standing waves.

4. Conclusion

The data presented in this short report demonstrates that in the period May to June 1979 the area between 5°N and 5°S at 92°E was subject to oscillating currents with a magnitude of 1 to 2 knots and a horizontal space scale of several degrees of latitude (400-500 km). The time scale of the oscillations is harder to estimate but appears to have been less than 2 weeks.

5. Acknowledgements

Thanks are due to the officers and men of HMAS DIAMANTINA for their help in the conduct of these experiments. Mr D. Rochford is thanked for some useful discussions and Dr. C. Nilsson for the use of his computer programmes for the dynamic height calculations for the RANRL transect.

6. References

Brown, N. and Hamon, B.V. (1961) 'An Inductive Salinometer' Deep-Sea Research 8, 65-75




Fig. 2(e). Temperature time series at 0°; 92°E on 16/5/79.




Fig. 3. Surface currents from ship's sets for (a) RANRL's south-going leg 12/5/79-15/5/79, with values at Equator on 16/5/79 and (b) CSIRO's south-going leg 28/5/79-1/6/79. (1cm \equiv 1 knot.)




Fig. 4. Dynamic height of sea - surface relative to 300m.




Fig. 4(d). Dynamic height relative to 1000db along 92°E.




Fig.5 Path of satellite-tracked buoys.

Cresswell, G.R., Richardson, G.T., Wood, J.E. and Watts, R. 'The CSIRO satellite-tracked torpedo buoy'. CSIRO Division of Fisheries Oceanography Report No 28, 1978.

Eriksen, C.C. (1979) 'An equatorial transect of the Indian Ocean'. J. Mar. Res. 37, 215-232.

O'Brien, J.J. and Hurlburt, H.E. (1974) 'Equatorial Jet in the Indian Ocean: Theory' Science, 184, 1075-1077.

Philander, S.G.H. (1973) 'Equatorial Undercurrent: Measurements and Theories' Reviews of Geophysics and Space Physics, 11, 513-570

Taft, B.A. and Knauss, J.A. (1967) 'The equatorial undercurrent of the Indian Ocean as observed by the Lusiad expedition' Bull. Scripps Inst. Oceanogr. 9.

Wyrtki, K. (1973) 'An Equatorial Jet in the Indian Ocean' Science, 181, 262-264